

Evaluation of Antibiotic Resistance and Susceptibility Patterns in Multidrug-Resistant *Acinetobacter* spp

Bahasht Yilmaz Ezzulddin ^{1*}

¹ M.Sc. Microbiology, Department of Microbiology, Azadi Teaching Hospital, Kirkuk, Iraq

E-mail: bahashtyilmaz@ntu.edu.iq

*Corresponding author

ARTICLE INFO

Keywords:


Acinetobacter species, VITEK 2, Antibiotic Resistance, Multidrug resistance, Imipenem.

Vol. 01, No. 05, 2026

<https://doi.org/10.5281/zenodo.23727637>

Copyright © 2026 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CCBY4.0).

<http://creativecommons.org/licenses/by/4.0/>



ABSTRACT

Antimicrobial-resistant *Acinetobacter* species (spp.) have emerged as significant healthcare-associated pathogens because they can develop resistance to many antibiotics, making infections difficult to treat. This study aims to evaluate antibiotic resistance and susceptibility patterns among *Acinetobacter* (A.) species. Six hundred twenty-four (624) clinical samples were collected from both inpatients and outpatients. Forty-four (44) *Acinetobacter* spp. isolates were identified by culture and biochemical characteristics and then confirmed with the VITEK 2 system. The antibiotic susceptibility profile was determined using the VITEK 2 AST-419 card. *Acinetobacter* spp. represented 7.1% of all isolates. The *Acinetobacter baumannii* complex was the most common, with 35 (79.6%), followed by *A. haemolyticus* 7 (15.9%) and *A. lwoffii* 2 (4.5%) isolates. The antibiotic susceptibility test showed the highest resistance rates to ampicillin/sulbactam 38 (86.4%), piperacillin 34 (77.3%), cefotaxime 29 (65.9%), levofloxacin 29 (65.9%), ciprofloxacin 28 (63.6%), ceftazidime 28 (63.6%), ticarcillin/clavulinate 27 (61.4%), and imipenem 27 (61.4%). Most isolates were sensitive to minocycline 34 (77.3%) and colistin 2 (4.5%), respectively. The data showed increased resistance to common antibiotics. Colistin, minocycline, and amikacin were effective antibiotic for managing infections induced by *Acinetobacter* species. Extensive drug resistance (XDR) and multidrug resistance (MDR) were the most prevalent among the *Acinetobacter* spp. isolates.

Introduction

Opportunistic infections are often associated with *Acinetobacter* spp., which are Gram-negative, nonfermentative bacteria that are present ubiquitously [1]. Human infections attributed to these bacteria involve both hospital-acquired and community-acquired infections, notably in critically ill patients with weakened host defenses. These infections include pneumonia, meningitis, and endocarditis, cutaneous and wound infections, peritonitis in patients on peritoneal dialysis, urinary tract infections (UTIs), and bacteremia [2]. The evolutionary timeline, phylogenetic diversity, and ecological prevalence are evident in the abundance of *Acinetobacter* species. Until September 2025, the genus had 87 species with validly published names, alongside several other taxonomically distinct provisional taxa and strains awaiting taxonomic clarification [3].

A multi-year hospital study revealed that *A. baumannii* was the most prevalent causative agent of *Acinetobacter* infections, followed by the non-*baumannii* species: *A. pittii* and *A. nosocomialis*; infections with *A. haemolyticus*, *A. johnsonii*, *A. junii*, *A. calcoaceticus*, *A. lwoffii*, *A. schindleri*, *A. ursingii*, and *A. courvalinii* have also been identified [4].

The emergence of Antimicrobial resistance has generated great concern, especially inside hospitals, as those receiving medical care or having compromised immune defenses are generally more susceptible to infections. Additionally, bacteria resistant to antibiotics can transmit within and across healthcare facilities, thereby resulting in diseases in patients termed healthcare-associated infections (HAIs), or they can spread to the community or environment (like soil and water) [5]. The significant frequency of resistant isolates among many antibiotic classes complicates initiation of successful empirical therapy. Owing to the variable multidrug resistance patterns exhibited by *Acinetobacter* spp., it is crucial to determine the susceptibility profiles common within the institution [6]. Numerous types

of acquired resistance mechanisms, including the production of extended-spectrum beta-lactamase enzymes (ESBLs), modification enzymes for aminoglycosides, modified binding sites for quinolones, and diverse efflux mechanisms, lead to significant challenges for clinicians when selecting an appropriate empirical antimicrobial agent. [7]. Consequently, in 2017, the World Health Organization (WHO) identified carbapenem-resistant *Acinetobacter baumannii* (CRAB) as a critical priority for the discovery and formulation of newly developed antibiotics [8]. This study aims to isolate and identify *Acinetobacter* spp. from various clinical samples, evaluate the antibiotic resistance and susceptibility patterns of *Acinetobacter* spp. (Multidrug-resistant, extensively drug-resistant, and pan-drug-resistant).

Materials and Methods

Study design, setting and patients

This cross-sectional study was conducted between February 2026 and June 2026 at Azadi Teaching Hospital in Kirkuk City, Iraq. Six hundred twenty-four (624) different clinical samples were obtained from both inpatients and outpatients, including urine, blood, sputum, burn swabs, wound swabs, high vaginal swabs, cerebrospinal fluid (C.S.F.), and bronchial wash. The criteria for inclusion in the study were suspected bacterial infections, whereas duplicate cultures and contaminated samples were excluded.

Sample collection, isolation, and bacteriological identification of *Acinetobacter* spp.

The samples collected from patients were classified as infected based on the clinical manifestations exhibited by each patient. All samples were collected by following strict aseptic protocols and were conveyed to the microbiology laboratory for processing.

The specimens were subsequently inoculated onto MacConkey agar, blood agar, and a medium, then incubated at 37°C for 24 hours. Bacterial isolates were identified by conventional microbiological methods based on colony morphology and preliminary testing, such as Gram staining, catalase, and oxidase assays. Traditional biochemical tests used, including Kligler iron agar (KIA), indole, citrate, urease, and motility assays (semi-solid manitol; SSM), to identify *Acinetobacter* species. Finally, all *Acinetobacter* species were confirmed using VITEK2 Gram-negative (GN) identification cards following the manufacturer's guidelines.

Antimicrobial susceptibility testing of *Acinetobacter* spp.

Testing for susceptibility to antimicrobial agents was carried out on all isolated *Acinetobacter* spp. using the VITEK 2 GN AST-N419 cards. The system-determined MIC values were measured according to CLSI 2026 standards [9], and the susceptibility level of each isolate was assessed as sensitive (S), intermediate (I), or resistant (R). According to the Centers for Disease Control and Prevention (CDC), isolates that are resistant to at least one antibiotic from three or more classes are defined as MDR. Isolates that demonstrate resistance to at least one antibiotic from each group of antibiotics are identified as XDR. Pan-drug-resistant (PDR) isolates are defined as strains that are resistant to all antibiotics evaluated in all groups of antimicrobial agents [10]. The identification of drug-resistant patterns (MDR, XDR, PDR, and Carbapenemase *Acinetobacter* spp. isolates) was obtained using VITEK 2 results, and percentages were determined in relation to the total isolates.

Statistical analysis

The data were analysed utilising SPSS software (version 27.0), and the chi-square (X^2) test was applied to assess the significance of percentages at probability levels of 0.05 and 0.01.

Results and Discussion

Patients and *Acinetobacter* spp. isolates characteristics

Six hundred twenty-four (624) clinical samples were obtained from patients at Azadi Teaching Hospital in Kirkuk City. Based on culture and biochemical testing, 190 (30.4%) of the 624 samples showed bacterial growth, including 44 (7.1%) *Acinetobacter* spp. The remaining 146 (23.3%) involved other bacteria, whereas 434 (69.6%) samples exhibited no bacterial growth, as listed in **Table (1)**.

Table (1): Culture results for different clinical samples.

Sample Culture		Number (No.)		Percentage (%)	
Bacterial Growth	Acinetobacter spp.	190	44	30.4%	7.1%
	Other bacteria		146		23.3%



No Bacterial Growth	434	69.6%
Total	624	100%

Our results were similar to those of the previous study [11], which reported that bacterial growth accounted for 28% in Thiruvallur City, India, and that the frequency of *Acinetobacter* spp. isolates was 9.9% among different clinical samples.

Furthermore, a previous study [12] reported a 4.9% frequency of *Acinetobacter* spp., agreeing with our findings. By contrast, our findings disagreed with a study carried out in Babylon Province [13], which found 20 (3.33%) *Acinetobacter* spp. isolates among 600 isolates from different clinical samples. The observed differences in frequency might be resulting from differences in study design, methodology, and timing [14].

The frequency of *Acinetobacter* spp. isolates in clinical samples was highest in burn swabs (29.6%), followed by urine (25%), blood (18.2%), sputum (13.6%), wound swabs (4.5%), high vaginal swabs (4.5%), cerebrospinal fluid (2.3%), and bronchial wash (2.3%), as shown in **Table (2)**.

Table (2): Isolation source and percentage of *Acinetobacter* species.

Isolation Source	Number of <i>Acinetobacter</i> spp.	Percentage of <i>Acinetobacter</i> spp.
Burn swabs	13	29.6%
Urine	11	25%
Blood	8	18.2%
Sputum	6	13.6%
Wound swabs	2	4.5%
Higher vaginal swabs	2	4.5%
Bronchial wash	1	2.3%
C.S.F.	1	2.3%
Total	44	100%

A comparable study carried out by [15] reported that the frequency of *Acinetobacter* spp. isolates was high in burn swabs (41%). Additionally, another study [16] conducted in Duhok Province reported that the frequency of *Acinetobacter baumannii* isolates was high in burn swabs (36.5%), consistent with our findings regarding the burn swabs prevalence rate. An increased number of isolates were observed in burn swabs due to the increased vulnerability of burns and wounds to contamination from the hospital environment, thus facilitating the spread of resistant strains. Additionally, most antibiotics demonstrate limited ability to penetrate the sites of burns and wounds [17].

The second most prevalent *Acinetobacter* spp. frequency in our study was observed in urine samples at a rate of 25%, which aligns with study [18], which reported a frequency of *Acinetobacter* spp. in urine of 25.43%. The high recovery rate from urine samples may reflect healthcare-associated urinary tract infections. *A. baumannii* is capable of causing infections at various clinical sites in individuals receiving hospital care. This pattern might reflect the frequent use of invasive medical devices, specifically urinary catheters [19].

Out of 624 patients, 352 were inpatients and 272 outpatients. The frequency of *Acinetobacter* spp. was higher in inpatients, 31 (70.5%), relative to outpatients, 13 (29.5%). Our findings revealed an increased prevalence of this infection among inpatients, which clearly reflects the nosocomial origin of this opportunistic pathogen. A similar occurrence was observed in other studies [20, 21]. It may last for extended periods in hostile environments (walls, surfaces, and medical equipment) within healthcare settings, and it is chromosomally resistant to many antibiotic classes [22].


There were 367 females and 257 males aged from less than one to more than 80 years. The females 30 (68.2%) appeared to be more infected than males 14 (31.8%). This finding agreed with study [23] conducted in Wasit, Iraq, which found that the infection rate in females was 38 (65.5%) more than in 20 (34.5%) males. This might be attributed to the higher frequency of hospital visits reported by females in comparison to males [20].

Regarding patient ages, the distribution of *Acinetobacter* spp. was highest in 16 patients (36.4%) aged 21–40 years whereas 10 patients (22.7%) were aged 61–80 years, 7 patients (15.9%) were aged 41–60 years, 5 patients (11.4%) were aged less than one year, 4 patients (9.1%) aged 1–20 years and 2 patient (4.5%) belonged to the age group more than 80 year.

Most infections occurred in age group 21–40 years (36.4%), and the lowest percentage was observed in the age group older than 80 years (4.5%); these results are in line with the prevalence reported by a recent local study [24] carried out in Babylon and Al Najaf City. The increasing frequency in the 21–40 age range could be linked to increased exposure to traumatic injuries, surgical interventions, and healthcare facilities, particularly in contexts where young and middle-aged individuals constitute a substantial portion of patients. A study [25] revealed a high distribution of *Acinetobacter* spp. among those aged over 50 years; our results align with this and showed a second high rate in the 61–80 age range, followed by the 41–60 group. This suggests that elderly individuals, particularly those with underlying comorbidities, are at greater risk for *A. baumannii* infections.

Isolation and bacteriological identification of *Acinetobacter* spp. isolates

The colonies morphology on MacConkey and blood agar were used for the preliminary identification of *Acinetobacter* spp. isolates (No. 44). The colonies of *Acinetobacter* spp. seen on MacConkey agar were round, pale-colored, and non-lactose fermenting. The colonies were observed to be grey or white with a convex morphology on blood agar, as shown in **Figure (1)**.



A.

MacConkey agar.

B. Blood agar.

Figure (1): Culture characteristics of *Acinetobacter* spp. on MacConkey and blood agar.

Biochemical tests, including oxidase (negative), catalase (positive), urease (variable), citrate utilization (positive), KIA (alkaline/alkaline), motility (non-motile), and indole (negative), were performed on the bacterial isolates as shown in **Figure (2)**.



Figure (2): Biochemical characterization of *Acinetobacter* spp. isolates.

In addition, the identification of 44 isolates of *Acinetobacter* species was confirmed by the VITEK 2 system, which differentiated them into three different species: *Acinetobacter baumannii* complex, *Acinetobacter lwoffii*, and *Acinetobacter haemolyticus*. The *A. baumannii* complex was found to be the most common, with 35 (79.6%), followed by *A. haemolyticus* 7 (15.9%) and *A. lwoffii* 2 (4.5%) isolates, which agreed with the findings of other studies [26] but disagreed in that they also isolated *A. radioresistens* and *A. junii*.

Antimicrobial Susceptibility Testing (AST)

The highest resistance pattern was detected against ampicillin/sulbactam 38 (86.4%), piperacillin 34 (77.3%), cefotaxime 29 (65.9%), levofloxacin 29 (65.9%), and ciprofloxacin 28 (63.6%). Additionally, resistance levels toward ceftazidime were 28 (63.6%), ticarcillin/clavulinate 27 (61.4%), and imipenem 27 (61.4%). An intermediate resistance pattern was observed toward cefepime 26 (59.1%), trimethoprim/ sulfamethoxazole 25 (56.8%), gentamicin 25 (56.8%), tobramycin 24 (54.5%), piperacillin/ tazobactam 24 (54.5%), and meropenem 23 (52.3%). The lowest resistance was observed against colistin 2 (4.5%) and minocycline 9 (20.4%), respectively, as shown in **Table (3)**.

Table (3): Antibiotic susceptibility tests of *Acinetobacter* species.

Class of antibiotics	Antibiotics	R	%	I	%	S	%	T
Penicillins	Piperacillin	34	77.3	2	4.5	8	18.2	44
B-lactam combination agents	Ampicillin/ Sulbactam	38	86.4	1	2.3	5	11.3	44
	Ticarcillin/Clavulinate	27	61.4	3	6.8	14	31.8	44
	Piperacillin/ Tazobactam	24	54.5	1	2.3	19	43.2	44
Cephalosporins	Ceftazidime	28	63.6	0	0	16	36.4	44
	Cefepime	26	59.1	6	13.6	12	27.3	44
	Cefotaxime	29	65.9	3	6.8	12	27.3	44
Carbapeneams	Imipenem	27	61.4	1	2.3	16	36.4	44
	Meropenem	23	52.3	2	4.5	19	43.2	44
Lipopeptide	Colistin	2	4.5	0	0	42	95.5	44
Aminoglycosides	Amikacin	16	36.4	8	18.2	20	45.4	44
	Gentamicin	25	56.8	1	2.3	18	40.9	44
	Tobramycin	24	54.5	0	0	20	45.5	44
Tetracycline	Minocycline	9	20.4	1	2.3	34	77.3	44
Fluoroquinolones	Ciprofloxacin	28	63.6	3	6.8	13	29.6	44
	levofloxacin	29	65.9	3	6.8	12	27.3	44
Folate pathway antagonists	Trimethoprim/ Sulfamethoxazole	25	56.8	0	0	19	43.2	44

* Total: T.

The resistance patterns to the selected antibiotics revealed a significant variation among *Acinetobacter* spp. isolates. The high resistance to piperacillin shown in our results aligns with study results [27], which reported a resistance rate of 88.8%.

A recent study [19] carried out in Baghdad City in 2026 reported high resistance to cefotaxime, with 30 (76.9%) showing resistance, followed by ceftazidime with 23 (59.0%) resistant. The resistance rates to piperacillin/tazobactam were 21 (53.8%), aligning with the findings of the current study. A variety of β -lactamases present in *Acinetobacter* species hydrolyze β -lactams and provide resistance to cephalosporins, carbapenems, and penicillins. Chromosome-encoded AmpC cephalosporinases confer resistance to broad-spectrum cephalosporins [27].

Our findings were in agreement with the study [28] in which the resistance rate toward Ciprofloxacin exhibited 77.7% resistance, followed by levofloxacin 66.6% and gentamicin 63.9%. Another study [17] showed that the tobramycin resistance level was 60.9%, which was similar to our findings. Lower sensitivity to fluoroquinolones further limits the accessibility of efficacious oral therapeutic alternatives for multidrug-resistant infections [29].

A study [30] carried out in Europe reported that, among the carbapenem class, imipenem had a resistance rate of 66%, comparable to our results, but disagreed in that they reported higher meropenem resistance (74%) than our findings. This increasing resistance is alarming, particularly as many patients previously had received treatment at primary or secondary care facilities with combinations of β -lactams, aminoglycosides, or fluoroquinolones, which contributed to selective pressure and resistance [31].

The low resistance patterns of amikacin demonstrate that they are effective therapeutically; a similar result was obtained from a retrospective study [32] carried out in India, in which the resistance level toward amikacin was 46%, indicating high aminoglycoside activity.

This study's results closely aligned with the study [33] carried out in Jordan that observed the level of resistance to ampicillin-sulbactam of 80% and trimethoprim/sulfamethoxazole of 56%, respectively.

Regarding minocycline, our result was similar to the study [21], which reported resistance to minocycline was 14.28%.

A local study [17] reported that the sensitivity rate of *A. baumannii* isolates to colistin was 97.5%, aligning with the results of our study. Colistin was the most effective antibiotic for managing infections induced by *Acinetobacter* species.

In this study 44 isolates of *Acinetobacter* spp., of these isolates, 9 (20.5%) were found to be MDR strains, 25 (56.8%) isolates XDR, 1 (2.2%) isolates PDR and 9 (20.5%) were non-MDR isolates, which was statistically highly significant (p -value < 0.001*) as mentioned in **Table (4)**. Antibiotic susceptibility testing performed by the VITEK-2 system indicated that 28 (63.6 %) of the *Acinetobacter* spp. isolated strains were carbapenemase producers phenotypically, as mentioned in **Table (5)**.

Table (4): Drug resistance patterns for *Acinetobacter* species.

Type of resistance	Number	Percentage	P-value
MDR	9	20.5	$p < 0.001^*$
XDR	25	56.8	
PDR	1	2.2	
Non-MDR isolates	9	20.5	
Total	44	100	

* $\chi^2 = 27.64$.

Table (5): Detection of carbapenemase-producing *Acinetobacter* species.

<i>Acinetobacter</i> species	Number	Percentage
CR- <i>Acinetobacter</i> spp.	28	63.6
Non -CR- <i>Acinetobacter</i> spp.	16	36.4
Total	44	100

Such resistance patterns have been observed in a local study [34] carried out in Diyala City, and [24] conducted in Babylon and Al-Najaf Cities, where *A. baumannii* isolates (38.0%) MDR and (62.0%) XDR strains reported were nearly comparable with the results of this study but disagreed in that there were also PDR and non-MDR isolates in our isolates. The significant proportion of MDR and XDR isolates identified in this study can be attributed to the exceptional adaptability of *A. baumannii*. Bacteria can develop many strategies for acquiring resistance to antimicrobial drugs, including the production of β -lactamases and carbapenemases, reducing outer membrane permeability, modifying targets, and enhancing the expression of multidrug efflux systems [19].

In a hospital-based prospective study [18] found that among carbapenem-resistant isolates screened (83.72%) confirmed the production of carbapenemase by the isolates which was higher percentage than our results. The observed variation may be attributed to elevated local antibiotic pressure, variations in methodology, the prevalence of resistance clones, and variations in geographical or institutional strategies for managing nosocomial infections [12]. Therefore, local monitoring studies of this nature play an important role in determining the most effective treatments for *Acinetobacter* infections.

CONCLUSION

Among *Acinetobacter* species, the *A. baumannii* complex appears to be isolated most frequently, followed by *A. haemolyticus*; the highest prevalence was found in burn swabs, followed by urine samples. Most *Acinetobacter* species were extensively drug-resistant, multidrug-resistant, and carbapenemase producers. This demonstrates the significance of alternative treatment approaches, stringent infection control protocols, and appropriate antibiotic use, especially in burn and medical wards.

REFERENCES

1. Ahmad S, Shakireen N, Khan MS, Mumtaz H, Ahmad W, Shah MH, Ahmad I, Khan W, Khan F, Nadeem A, Naqvi N. Prevalence and antimicrobial susceptibility of *Acinetobacter* spp. in a tertiary care hospital in Peshawar: a cross-sectional study. *Ann Med Surg (Lond)*. 2023;85(5):1584–1589.
2. Mohammed SH, Ahmed MM, Alameer AA, Abd Alabbas HH, Mohammad AZ, Abed AW, Mohsin ZA, Jasim MZ, Abd Al Hamza MZ, Abid ZN. Prevalence of *Acinetobacter* species isolated from clinical samples referred to Al-Kafeel Hospital, Iraq and their antibiotic susceptibility patterns from 2017–2021. *Iran J Med Microbiol*. 2022;16(1):76–82.
3. Karah N, Nemec A, Uhlin BE. History of the taxonomy of *Acinetobacter*: the emergence of hospital-adapted species of global health concern. *Int J Syst Evol Microbiol* 2025;75(12):1-26.
4. Wimalasekara RL, White D, Kumar A. Targeting *Acinetobacter baumannii* resistance-nodulation-division efflux pump transcriptional regulators to combat antimicrobial resistance. *npj Antimicrob Resist*. 2025;3(1):4.
5. Silva ME, Gomes MA, Rodrigues RS, Lima NC, Carvalho AG, Taborda RL, Matos NB. Multidrug-resistant *Acinetobacter* spp. from hospital intensive care units in Brazilian Amazon. *Braz J Infect Dis*. 2023;27(6):103687.
6. Singh BK, Darryl G, Kumari S, Bagul K. Prevalence of *Acinetobacter* spp. in lower respiratory tract infections and its antimicrobial susceptibility pattern in a tertiary care hospital. *Int J Res Med Sci*. 2026;14(5):1922-1926.
7. Tewari R, Chopra D, Wazahat R, Dhingra S, Dudeja M. Antimicrobial susceptibility patterns of an emerging multidrug resistant nosocomial pathogen: *Acinetobacter baumannii*. *Malays J Med Sci*. 2018;25(3):129–134.
8. Müller C, Reuter S, Wille J, Xanthopoulou K, Stefanik D, Grundmann H, Higgins PG, Seifert H. A global view on carbapenem-resistant *Acinetobacter baumannii*. *MBio*. 2023;14(6):e02260-23.
9. CLSI, Performance Standards for Antimicrobial Susceptibility Testing, 36th ed. (Clinical and Laboratory Standards Institute, 2026): CLSI supplement M100.
10. Mahto M, Chaudhary M, Shah A, Show KL, Moses FL, Stewart AG. High antibiotic resistance and mortality with *Acinetobacter* species in a tertiary hospital, Nepal. *PHA*. 2021;11(s1):13-17.
11. Vasumathi A, Suganthi M, Deepa R, Thilakavathi N. Isolation, identification and antimicrobial susceptibility profile of *Acinetobacter baumannii* from various clinical specimens received at a tertiary care hospital. *Int J Acad Med Pharm*. 2024;6(4):1086-1088.
12. Tchinda CF, Djim-Adjim-Ngana K, Mbogning TA, Simo BF, Baira SB, Feudjieu GE, Matchuenkam SG, Nfor GN, Takuissu GR, Essola KJ, Walyaro CC. Prevalence and risk factors associated with infections linked to carbapenemase-producing *Acinetobacter* species circulating in the city of Yaoundé, Cameroon. *IJID regions*. 2025;16:100698.
13. Al-Hasnawy HH, Rahi AA, Hadi BH. Molecular Characterization of bla OXA-51 Gene among *Acinetobacter baumannii* Isolates in Babylon Province. *Hilla Univ Coll J Med Sc*. 2024;2(2):24-32.
14. Mohammed SH, Ahmed MM, Alameer AA, Abd Alabbas HH, Mohammad AZ, Abed AW, Mohsin ZA, Jasim MZ, Abd Al Hamza MZ, Abid ZN. Prevalence of *Acinetobacter* spp. isolated from clinical samples referred to Al-Kafeel Hospital and their antibiotic susceptibility patterns from 2017-2021. *Iran J Med Microbiol*. 2022;16(1):76-82
15. Jaber AK, Hasan AM, Kkudhaier SR. Phenotypic and genotypic analysis of antibiotic resistance in *Acinetobacter* spp. isolates in Baghdad, Iraq. *Microbes Infect Dis*. 2025.

16. Haji MN, Al-Brefkanih AMT. Detection and characterization of carbapenem resistant *Acinetobacter baumannii* isolated from different clinical specimens in Duhok Province–Iraq. *Hist Med*. 2023;9(1):783–794.
17. Sehree MM, Abdullah HN, Jasim AM. Isolation and evaluation of clinically important *Acinetobacter baumannii* from intensive care unit samples. *J Tech*. 2021;3(3):83-90.
18. Rahman MM. Detection of blaOXA-23 gene among carbapenem-resistant *Acinetobacter* species isolated from various clinical samples. *Eur J Mol Clin Med*. 2022;9(03):9803-9810.
19. Moshin AS, Jamil HA. Prevalence of multidrug-resistant and carbapenem-resistant *Acinetobacter baumannii* among clinical specimens in Baghdad, Iraq. *Shifaa*. 2026;36-44.
20. Rebic V, Masic N, Teskeredzic S, Aljicevic M, Abduzaimovic A, Rebic D. The importance of *Acinetobacter* species in the hospital environment. *Med Arch*. 2018;72(5):325–329.
21. Rani P, Payal N, Khandait M, Dixit S. Identification and determination of the AST pattern of *Acinetobacter* species isolated from different clinical samples by VITEK[®] Compact. *medRxiv*. 2026.
22. Sharmin N, Khoda MM, Uddin MN, Shamsuzzaman SM, Jahan H. Prevalence, antibiotic resistant pattern and genotypic detection of *Acinetobacter baumannii* isolated from different clinical specimens of patients admitted at a tertiary care hospital in Bangladesh. *Bangladesh J Med Microbiol*. 2024;18(1):37-43.
23. Alawi AJ, Pourahmad F, Fattahi K. Detection of carbapenemase genes in *Acinetobacter baumannii* isolates from Al-Azizya Hospital, Wasit, Iraq. *Jundishapur J Microbiol*. 2025;18(6):1-9.
24. Kadhim AA, Hadi ZJ. High prevalence of bla NDM and bla OXA carbapenemases in multiple antibiotic resistant *Acinetobacter baumannii* from Iraqi Hospital. *Kufa Med J*. 2025;21(2):30.
25. Dhanapal B, Risha M, Saikumar C. Emerging drug resistance in *Acinetobacter* species: a study on isolation, speciation, and antimicrobial susceptibility patterns in a tertiary care hospital. *Biomed Pharmacol J*. 2025;18(2):1591-1599.
26. Kumari M, Bhattarai NR, Rai K, Pandit TK, Khanal B. Multidrug-resistant *Acinetobacter*: detection of ESBL, MBL, blaNDM-1 genotype, and biofilm formation at a tertiary care hospital in Eastern Nepal. *Int J Microbiol*. 2022;1(1):1-9.
27. Khalaf SA, Mohammed MR, Sabah ZA. Prevalence of *Acinetobacter baumannii* in Diyala Governorate from different clinical sources and its antibiotics profile analysis. *Int J Trop Dis Health*. 2024;45(9):63-70.
28. Hamid AA, Al-Lami SM, Saber ZM. Molecular characterization of OXA carbapenemase genes in *Acinetobacter baumannii* isolated from different clinical samples. *Cell Physiol Biochem*. 2026;60:445-457.
29. Kumar R, Maheshwari RK, Viridi SL, Parkash R. Isolation and antimicrobial susceptibility patterns of *Acinetobacter baumannii* with special emphasis on multidrug resistance. *IP Int J Med Microbiol Trop Dis*. 2026;12(3):552-556.
30. Müller H, Romano L. Detection and species-level identification of *Acinetobacter* spp. in clinical specimens and antimicrobial resistance profiles: A multicenter cross-sectional study in Europe. *Med Res Pract*. 2025;2(6):79-84.
31. Mahajan A. A study on isolated *Acinetobacter* species from various clinical samples of tertiary care centre, their identification and antimicrobial susceptibility pattern. *Int J Life Sci Biotechnol Pharma Res*. 2025;14(11):44-49.
32. Saikia P, Shetty K. Prevalence of *Acinetobacter* spp. and antibiotic susceptibility pattern in various clinical samples in a tertiary care hospital. *SSR Inst Int J Life Sci*. 2026;12(1): 9015-9023.
33. Gharaibeh MH, Abandeh YM, Elnasser ZA, Lafi SQ, Obeidat HM, Khanfar MA. Multi-drug resistant *Acinetobacter baumannii*: phenotypic and genotypic resistance profiles and the associated risk factors in teaching hospital in Jordan. *J Infect Public Health*. 2024;17(4):543-550.
34. Al-Masoodi A, Bdioui A, Alzubaidi A. Prevalence and resistance patterns of multidrug-resistant *Acinetobacter baumannii* in clinical samples from Diyala, Iraq. *Trop J Pharm Res*. 2025;24(12):1553-1562.