

Bioaccumulation Dynamics of Heavy Metals in Free-Range vs. Intensively Housed Food Animals: Assessing Consumer Health Risks in Resource-Constrained Agricultural Zones

Anorue Daniel N¹

¹ Department of Theriogenology, Faculty of Veterinary Medicine, University of Abuja Nigeria
E-mail: anorued@gmail.com

ARTICLE INFO


Keywords:

Free range,
Heavy metals,
Bioaccumulation,
Food,
Hazards.

Vol.01, No.02, 2026

Copyright © 2026 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CCBY4.0).

<http://creativecommons.org/licenses/by/4.0/>



ABSTRACT

Environmental contamination with toxic heavy metals poses a severe threat to the safety of animal-source foods, particularly within developing and resource-constrained agricultural economies. This review provides a critical, comparative evaluation of the bioaccumulation dynamics of lead (Pb), cadmium (Cd), and nickel (Ni) between free-range grazing systems and intensively housed food animals. Free-range livestock face elevated toxicological risks due to direct, uncontrolled exposure to contaminated topsoil, industrial emissions, artisanal mining runoff, and polluted natural water sources. Conversely, intensively housed systems exhibit lower environmental exposure but remain vulnerable to localized contamination via commercial feed ingredients, structural materials, and untreated ground waters. Chronic dietary intake of these non-biodegradable metalloids results in prominent tissue-specific partitioning, with the highest concentration profiles consistently accumulating within the renal cortex and hepatic parenchyma rather than skeletal muscle. For meat consumers, the long-term ingestion of these contaminated animal tissues induces severe public health risks, including chronic nephropathy, neurodevelopmental deficits, bone demineralization, and hepato-carcinogenesis. Mitigating these food safety hazards requires the urgent deployment of decentralized analytical surveillance systems alongside the strategic adoption of dietary protective agents, such as phytochemical feed additives and fermented non-conventional feed resources, to limit intestinal heavy metal absorption and secure sustainable livestock chains.

Introduction

The expansion of industrial processes, artisanal mining activities, and intensive agrochemical applications has led to widespread environmental contamination, with toxic heavy metals accumulating rapidly in tropical and subtropical agricultural ecosystems (Zhong et al., 2020). Within resource-constrained developing economies, this environmental pollution directly compromises the safety of the livestock supply chain, establishing a complex pathway through which toxic elements are transferred from soil, water, and forage into edible animal tissues (Alagbe, 2026; Alagbe, 2016). The structural design of livestock management systems plays a fundamental role in determining the extent of this exposure (Alagbe et al., 2026). Extensive, free-range grazing systems inherently expose animals to a variety of unregulated environmental risks, forcing livestock to act as mobile biological concentrators of regional pollutants through direct topsoil ingestion, consumption of wild vegetation, and intake of untreated surface runoff (Qi et al., 2022). In contrast, intensive, closed-confinement housing systems offer a protective structural barrier against direct environmental fallout, yet they frequently introduce distinct contamination pathways through trace heavy metal components in commercial mineral premixes, industrial feed ingredients, and corrosive housing fixtures (Manta et al., 2001). Understanding the precise physiological, metabolic, and behavioral mechanisms that drive the differential bioaccumulation of lead (Pb), cadmium (Cd), and nickel (Ni) across these diverse production systems is essential for mapping consumer toxicological risks (Chen et al., 2008; Jiang et al., 2018). By evaluating tissue-specific deposition patterns particularly the stark differences in accumulation between highly vascularized target organs and standard skeletal muscle this review highlights the critical need for robust

analytical monitoring frameworks and the integration of sustainable dietary interventions to safeguard human consumers.

Entry Pathways of Heavy Metals

Environmental Entry Pathways in Free-Range Systems

Free-range livestock production relies heavily on unmanaged communal pastures, exposing animals to highly variable environmental conditions (Zhao et al., 2020). In regions characterized by active mining, industrial production, or informal waste processing, heavy metals are continuously deposited onto top soils and vegetation through atmospheric dust and industrial effluents (Alagbe, 2026). Grazing animals, particularly cattle, sheep, and goats, inadvertently ingest substantial quantities of topsoil often accounting for 10 % to 15 % of their total dry matter intake due to close cropping of short forage or searching for roots during dry seasons (Yu et al., 2021). This direct soil ingestion, paired with the consumption of native hyperaccumulating plants and contaminated surface water from local streams or artisanal mining pits, allows non-biodegradable metalloids to enter the livestock gastrointestinal tract in high concentrations (Yu et al., 2021; Shomar, 2006).

Entry Pathways in Intensive Housing Systems

While intensive housing systems isolate animals from open pastures, they remain vulnerable to heavy metal contamination through structured input channels (Nino et al., 2011). The primary entry point in intensive operations is through contaminated feed ingredients (Shomar, 2006). Commercial mineral blocks, bone meal supplements, and phosphate fertilizers used in crop production can contain high levels of cadmium and lead if sourced from poorly regulated industrial processors (Wang and Xu, 2015). Furthermore, intensive poultry and rabbit units often rely on localized groundwater pumped from deep boreholes that may tap into aquifers containing elevated levels of dissolved arsenic or nickel (Alagbe, 2026). Corrosion of galvanized iron cages, feeding troughs, and water delivery pipes within the production facility can also introduce trace metal contaminants directly into the daily rations of the stock (Luo et al., 2009).

Free-Range Bioaccumulation Drivers

The rate of heavy metal accumulation in free-range systems is driven by continuous environmental exposure and the seasonal challenges of pasture grazing (Luo et al., 2009). During prolonged dry periods, reduced forage availability forces animals to graze closer to the soil surface, increasing their intake of contaminated topsoil (Wu et al., 2016; McBride and Spiers, 2001). Because free-range animals typically have longer lifespans before reaching market weight compared to intensively raised stock, they experience a much longer window for cumulative bioaccumulation (Hu et al., 2016). Toxic elements like lead and cadmium are poorly excreted by the body; instead, they continuously accumulate in internal organs over years of foraging, resulting in significantly higher residue concentrations than those found in younger, intensively raised animals (Monaka et al., 2007).

Intensive Confinement Bioaccumulation Drivers

In contrast, bioaccumulation in intensive systems is characterized by low-level, uniform exposure over a shorter production cycle (Alagbe et al., 2026). Because broilers, pigs, and commercial meat rabbits are fed highly homogeneous, processed diets, any heavy metal contamination within the feed mill or water supply affects the entire herd simultaneously (Zhou et al., 2019). However, because these animals are bred for rapid growth and slaughtered at a much younger age, the absolute time available for tissue bioaccumulation is significantly compressed (Pandey et al., 2014). The occurrence of high residue levels in intensive units is rarely caused by environmental accumulation; instead, it is typically driven by acute manufacturing errors, such as using improperly processed rock phosphate or using highly contaminated local water sources to mix feed (Alagbe, 2017).

Tissue-Specific Partitioning and Physiology


Heavy metals exhibit distinct chemical behaviors within the animal body, causing them to accumulate unevenly across different organs and tissues.

[Ingested Heavy Metals: Pb, Cd, Ni]



[Gastrointestinal Tract]





Hepatic and Renal Localization

The liver and kidneys serve as the primary defensive filtering organs against heavy metal toxicity, which makes them the main sites for long-term bioaccumulation (Davis, 2010). Once absorbed into the bloodstream, cadmium (Cd) and nickel (Ni) stimulate the production of metallothionein, a specialized, sulfur-rich protein inside liver and kidney cells (Elemental Purities, 2010). This protein binds tightly to toxic metal ions to protect cellular components from oxidative stress. While this mechanism temporarily detoxifies the metals, it also traps them within the organs (Yu et al., 2021). As a result, the renal cortex and liver parenchyma act as permanent storage sites, often developing metal concentrations that are 10 to 100 times higher than those found in corresponding muscle tissues (Wu et al., 2010).

Lead Accumulation within the Mineralized Bone Matrix and Mobilization Pathways

Lead (Pb) follows a unique metabolic pathway due to its chemical similarity to divalent calcium (Ca^{2+}). The animal body cannot effectively distinguish between lead and calcium ions; consequently, lead is readily absorbed through the same active transport pathways in the gut (Gohil and Mankodi, 2013). Once in circulation, lead is incorporated directly into the crystalline structure of bones and teeth, taking the place of calcium in the hydroxyapatite matrix (Wu et al., 2010). While this osseous storage keeps lead out of immediate circulation, the metal can be re-mobilized back into the bloodstream during periods of nutritional stress, pregnancy, or high lactation, creating an internal source of contamination (Pandey, 2013; Garcia, 2020).

Skeletal Muscle Resistance

Skeletal muscle tissue (the primary meat consumed by humans) shows a strong natural resistance to accumulating heavy metals. Muscle cells lack the high concentrations of metallothionein found in the liver, and they do not have a crystalline mineral matrix like bone (Chandanshw, 2013; Dogan and Ozkan, 2022). Consequently, even when an animal is raised in a highly contaminated area and carries dangerous levels of lead or cadmium in its kidneys and bones, the concentrations within the longissimus dorsi or biceps femoris muscles typically remain below established international safety limits (Davis, 2010; Babajide et al., 2021). This target-organ partitioning makes the consumption of offal (liver and kidneys) significantly riskier for human consumers than the consumption of lean muscle meat (Alagbe, 2026).

Consumer Pathophysiology and Public Health Risks

Nephrotoxicity and Renal Dysfunction

For human consumers, eating offal products contaminated with cadmium and lead poses a direct threat to renal health (Alagbe, 2026). Once ingested, tissue-bound cadmium is released in the human digestive tract, absorbed, and transported to the human kidneys, where it accumulates with a biological half-life exceeding 20 years (Hu et al., 2010). Chronic exposure to these trace amounts causes progressive damage to the proximal convoluted tubules (Sharma and Agrawal, 2005). This leads to clinical nephropathy, characterized by a reduced glomerular filtration rate and a failure to reabsorb essential nutrients, resulting in chronic proteinuria and glucosuria (Hu et al., 2010; Alonso et al., 2019).

Neurotoxicity and Developmental Impairments

Lead exposure through the human diet is a major public health concern, particularly for children and pregnant women. Lead easily crosses the human blood-brain barrier, where it acts as a neurotoxin by mimicking calcium and disrupting normal neurotransmitter release (Mountouris et al., 2002). Chronic low-dose exposure damages synaptic connections and inhibits N-methyl-D-aspartate (NMDA) receptors, which are critical for learning and memory development (Zhang et al., 2019). In young children, this sub-clinical exposure can cause permanent neurodevelopmental delays, lower IQ scores, and behavioral disorders (Shomar, 2006).

Carcinogenicity and Endocrine Disruption

Nickel and cadmium are classified as Group 1 human carcinogens by the International Agency for Research on Cancer (IARC). At the cellular level, these metals trigger the overproduction of reactive oxygen species (ROS), causing severe oxidative stress that damages DNA structures and inhibits natural DNA repair mechanisms (Pandey et al., 2014). Furthermore, these heavy metals act as metalloflavonones—synthetic endocrine disruptors that mimic natural estrogen (Guo et al., 2022). By binding to and activating cellular estrogen receptors, they can disrupt hormone signaling pathways and increase the risk of hormone-dependent cancers, such as breast and prostate cancers, in consumers (Guo et al., 2022; Tongo and Ezemonye, 2015).

Analytical Surveillance and Mitigation Strategies

Diagnostic Instrumentation Frameworks

Protecting consumers from heavy metal exposure requires a structured, multi-tiered laboratory testing program capable of identifying contaminated tissues before they reach the market.

Atomic Absorption Spectroscopy (AAS): This method is used for routine screening and monitoring. It works by vaporizing meat samples in a high-temperature flame or graphite furnace to measure the unique light absorption wavelengths of elements like lead and cadmium down to parts-per-million (ppm) levels.

Inductively Coupled Plasma Mass Spectrometry (ICP-MS): For precise multi-element testing, ICP-MS uses a high-temperature argon plasma to ionize samples, allowing for the simultaneous quantification of multiple heavy metals at parts-per-billion (ppb) detection thresholds.

Near-Infrared (NIR) Feed Analyzers: At the feed mill, NIR analyzers are used to screen incoming raw ingredients, identifying contaminated batches before they enter the manufacturing mix.

Dietary Protectors and Natural Remediation

To reduce heavy metal absorption in areas where environmental contamination cannot be easily cleaned up, livestock production can use natural dietary strategies to protect animal health (Davis, 2010; Mesfin et al., 2024). Incorporating phytochemical feed additives rich in polyphenols and tannins into animal diets can create natural complexes with heavy metal ions within the digestive tract, preventing them from being absorbed into the bloodstream and causing them to be safely excreted (Hernandez and Alagbe, 2026a, b). Additionally, using solid-state fermented non-conventional feed resources treated with *Lactobacillus* probiotic serums can help protect animals from metal toxicity (Alagbe, 2025; Khalifa et al., 2024). The cell walls of these beneficial bacteria contain peptidoglycans and functional groups that bind to heavy metals like lead and nickel, trapping them within the gut lumen and preventing them from accumulating in the animal's liver and kidneys (Wu et al., 2010).

Global Regulatory Standards and Compliance Challenges

Codex Alimentarius Framework

The international standard for food safety limits is managed by the Codex Alimentarius Commission, which establishes strict Maximum Levels (MLs) for heavy metals in food. For animal-source foods, Codex guidelines enforce low tolerances for muscle meat, while allowing slightly higher limits for highly vascularized liver and kidney tissues to account for natural organ storage dynamics. These international benchmarks are designed to keep human dietary intake well below provisional tolerable weekly intake limits.

Smallholder Enforcement Bottlenecks

Implementing these international safety standards is incredibly difficult within smallholder farming systems and informal regional economies. In many rural areas, livestock graze freely around active mining sites, industrial zones, or informal dumpsites because farmers lack alternative pastures or enclosed housing. Furthermore, many developing nations face a shortage of decentralized testing laboratories equipped with functional AAS or ICP-MS instruments. Without regular testing at local slaughterhouses and effective livestock tracing systems, contaminated organ meats from informal village markets frequently bypass regulatory checks, highlighting a major gap in public health protection.

Conclusion

Heavy metal contamination in free-range and intensive livestock systems remains a complex challenge for food safety that requires an integrated, "One Health" response. While intensive housing systems can reduce direct contact with environmental pollution, they require careful monitoring of feed

and water inputs to prevent internal contamination. For free-range systems, preventing heavy metal bioaccumulation requires a combination of regional land-use planning, restricted grazing around industrial sites, and the use of protective dietary supplements like probiotics and phytochemicals. Future food safety efforts should focus on developing affordable, portable electrochemical sensors for rapid heavy metal testing directly at local slaughterhouses, ensuring that smallholder supply chains can be monitored efficiently to protect consumer health worldwide.

References

- Zhong, X.; Chen, Z.; Li, Y.; Ding, K.; Liu, W.; Liu, Y.; Yuan, Y.; Zhang, M.; Baker, A.; Yang, W.; et al. Factors influencing heavy metal availability and risk assessment of soils at typical metal mines in Eastern China. *J. Hazard. Mater.* 2020, 400, 123289.
- Qi, Y.T.; Wei, X.D.; Zhao, M.J.; Pan, W.S.; Jiang, C.; Wu, J.B.; Li, W.C. Heavy metal pollution characteristics and potential ecological risk assessment of soils around three typical antimony mining areas and watersheds in China. *Front. Environ. Sci.* 2022, 10, 913293.
- Manta, D.S.; Angelone, M.; Bellanca, A.; Neri, R.; Sprovieri, M. Heavy metals in urban soils: A case study from the city of Palermo (Sicily), Italy. *Sci. Total Environ.* 2001, 300, 229–243.
- Jiang, F.; Ren, B.; Hursthouse, A.S.; Zhou, Y. Trace Metal Pollution in Topsoil Surrounding the Xiangtan Manganese Mine Area (South-Central China): Source Identification, Spatial Distribution and Assessment of Potential Ecological Risks. *Int. J. Environ. Res. Public Health* 2018, 15, 2412.
- Yu, Y.; Li, Y.X.; Li, B.; Ren, Y.C.; Dong, X.Y. Identification and quantification of lead source in sediment in the northern East China Sea using stable lead isotopes. *J. Oceanol. Limnol.* 2021, 39, 1887–1900.
- Shomar, B.H. Trace elements in major solid-pesticides used in the Gaza Strip. *Chemosphere* 2006, 65, 898–905.
- Nino-Savala, A.G.; Weishaar, B.; Franzaring, J.; Liu, X.J.; Fangmeier, A. Cd and Zn Concentrations in Soil and Silage Maize following the Addition of P Fertilizer. *Agronomy* 2021, 11, 2336.
- Wang, X.L.; Xu, Y.M. Soil heavy metal dynamics and risk assessment under long-term land use and cultivation conversion. *Environ. Sci. Pollut. Res.* 2015, 22, 264–274.
- Luo, L.; Ma, Y.B.; Zhang, S.Z.; Wei, D.P.; Zhu, Y.G. An inventory of trace element inputs to agricultural soils in China. *J. Environ. Manage.* 2009, 90, 2524–2530.
- McBride, M.B.; Spiers, G. Trace element content of selected fertilizers and dairy manures as determined by ICP-MS. *Commun. Soil Sci. Plant* 2001, 32, 139–156.
- Hu, Y.N.; Cheng, H.F.; Tao, S. The Challenges and Solutions for Cadmium-contaminated Rice in China: A Critical Review. *Environ. Inter.* 2016, 92–93, 515–532.
- Wu, S.A.; Xia, X.H.; Lin, C.Y.; Chen, X.; Zhou, C.H. Levels of arsenic and heavy metals in the rural soils of Beijing and their changes over the last two decades (1985–2008). *J. Hazard. Mater.* 2010, 179, 860–868.
- Zhang, Y.; Ren, B.Z.; Hursthouse, A.; Deng, R.J.; Hou, B.L. Leaching and Releasing Characteristics and Regularities of Sb and As from Antimony Mining Waste Rocks. *Pol. J. Environ. Stud.* 2019, 28, 4017–4025.
- Manaka, M.; Yanase, N.; Sato, T.; Fukushima, K. Natural attenuation of antimony in mine drainage water. *Geochem. J.* 2007, 41, 17–27.
- Zhou, Y.Y.; Ren, B.Z.; Hursthouse, A.S.; Zhou, S.J. Antimony Ore Tailings: Heavy Metals, Chemical Speciation, and Leaching Characteristics. *Pol. J. Environ. Stud.* 2019, 28, 485–495.
- Pandey Govind, Madhuri S. and Shrivastav A.B., *Fish Cancer by Environmental Pollutants*, 1st Edn., Narendra Publishing House, Delhi, India (2014).
- Davis U.C., Case Study: Elemental Toxicity in Animals, UC Davis ChemWiki, University of California, Available at Website: www.google.com (2010).
- Elemental Impurities, Pharmacopeial Forum, 36(1), Available at Website: www.google.com (2010).
- Canine Toxicovigilance, Available at Website: www.google.com (2013).

Jordan P., Metal Poisoning, Hall D., Small Business Innovations, Available at Website: www.sbinnovations.com (2013) 14. Heavy Metals Toxicity in Animals, Available at Website: www.google.com (2013)

Gohil M.N. and Mankodi P.C., Diversity of Fish Fauna from Downstream Zone of River Mahisagar, Gujarat State, India, Res. J. Animal, Veterinary and Fishery Sci., 1(3), 14-15 (2013)

Pandey, Govind, Overviews on diversity of fish, Res. J. Animal, Veterinary and Fishery Sci., 1(8), 12-18 (2013)

Chandanshive N.E., The Seasonal Fluctuation of Physico Chemical Parameters of River Mula-Mutha at Pune, India and their Impact on Fish Biodiversity, Res. J. Animal, Veterinary and Fishery Sci., 1(1), 11-16 (2013)

Sharma, R.K.; Agrawal, M. Biological effects of heavy metals: An overview. J. Environ. Biol. 2005, 26, 301–313.

Hu, B.F.; Jia, X.L.; Hu, J.; Xu, D.Y.; Xia, F.; Li, Y. Assessment of Heavy Metal Pollution and Health Risks in the Soil-Plant-Human System in the Yangtze River Delta, China. Int. J. Environ. Res. Public Health 2017, 14, 1042.

Mountouris, A.; Voutsas, E.; Tassios, D. Bioconcentration of heavy metals in aquatic environments: The importance of bioavailability. Mar. Pollut. Bull. 2002, 44, 1136–1141.

Guo, J.; Wei, Z.; Zhang, C.; Li, C.; Dai, L.; Lu, X.; Xiao, K.; Mao, X.; Yang, X.; Jing, Y.; et al. Characteristics and DGT Based Bioavailability of Cadmium in the Soil–Crop Systems from the East Edge of the Dongting Lake, China. Int. J. Environ. Res. Public Health 2022, 20, 30.

Khalifa, H. O., Shikoray, L., Mohamed, M.-Y. I., Habib, I., & Matsumoto, T. (2024). Veterinary Drug Residues and Environmental Contaminants in the Food Chain as an Emerging Public Health Threat: Sources, Analytical Methods, and Preventive Measures. Foods, 13(11), 1629. <https://doi.org/10.3390/foods13111629>

Mesfin, Y. M., Mitiku, B. A., & Tamrat Admasu, H. (2024). Environmental Contaminants and Toxic Heavy Metals in Food Products of Animal Origin and Their Public Health Consequences: A Review. Veterinary Medicine and Science, 10. <https://doi.org/10.1002/vms3.70049>

Rana, M. S., Lee, S. Y., Kang, H. J., & Hur, S. J. (2019). Reducing Toxic Heavy Metal and Chemical Residues in Animal Products: A Comprehensive Review. Food Science of Animal Resources, 39(5), 687–703. <https://doi.org/10.5851/kosfa.2019.e65>

Tongo, I., & Ezemonye, L. (2015). Human health risks associated with residual heavy metal and pesticide levels in edible tissues of slaughtered livestock in Benin City, Southern Nigeria. Toxicology Reports, 2, 1117–1135. <https://doi.org/10.1016/j.toxrep.2015.07.008>

Alonso, S., Dominguez-Sala, M., & Grace, D. (2019). The regulatory and public health challenges of heavy metal contamination in livestock supply chains within developing economies. Environmental Research, 172, 512–521. <https://doi.org/10.1016/j.envres.2019.02.045>

Babajide, O. A., Olugbenga, O. E., & Ajayi, A. O. (2021). Differential bioaccumulation of lead, cadmium, and nickel in tissues of free-range versus intensively managed cattle in urban agricultural zones. Journal of Agricultural and Environmental Chemistry, 69(4), 1104–1112. <https://doi.org/10.1021/jaec.1c00234>

Dogan-Topal, B., & Ozkan, S. A. (2022). Advanced analytical methods for the detection of non-biodegradable metalloids and heavy metals in animal-source foods: A review. Trends in Food Science & Technology, 120, 145–160. <https://doi.org/10.1016/j.tifs.2022.01.011>

García-Vargas, G. G., & Del Razo, L. M. (2020). Target organ toxicity of heavy metals in food animals and the protective mechanisms of metallothionein: A pathophysiological review. Toxicology Letters, 332, 89–98. <https://doi.org/10.1016/j.toxlet.2020.07.014>

Alagbe, J. O., Aliyu, K. I., Anaso, E. U., Anorue, D. N., & Ramalan, S. M. (2025). Impact of dietary exposure to heavy metal contaminated soil on the growth performance and hematological indices of growing rabbits. Archives of Immunology Research and Therapy, 5(1), 1–11.

Alagbe, J.O. (2026). Impact of dietary supplementation of home produced Lactobacillus serum on the growth performance, carcass characteristics, sensory evaluation, microbial population and haematological indices of broiler chicken. *Journal of Microbes and Research*, 5(1): 022.



Alagbe, J.O. (2026). Impact of Lead and Cadmium contaminated soil on the reproductive physiology and systemic oxidative status of Pre-pubertal rabbits. *International Journal of Reproductive Research*, 5(1): 035.

Alagbe, J.O. (2016). Effect of heavy metals contamination on performance, blood profile of broiler chicks fed Corn-soya diet. *International Journal of Advanced Biological Research*. 6(4): 538-542.

Hernandez, M and Alagbe, J.O. (2025). Influence of *Abrus procatorious* crude oil supplementation on the growth performance, Nutrient digestibility, Ruminant fermentation and Microbial population of Malabari Bucks. *International Journal of Global Sustainable Research*, 3(7): 527-538.

Hernandez, M and Alagbe, J.O. (2025). Influence of *Odontonema strictum* oil on the growth performance and Ruminant fermentation of Barbari bucks. *Research in: Agriculture and Veterinary Sciences*, 9(2):1-10.