

Chemical Residues in Intensified Livestock Production: A Review of Emerging Public Health Concerns and Mitigation Pathways

Alagbe, John Olujimi ¹

¹ Department of Animal Nutrition and Biochemistry, Gandhi College of Agriculture, Rajasthan, India
Department of Animal Science, Centre for Distance Learning and Continuing Education, University of Abuja,
Gwagwalada, Nigeria

E-mail: dralagbe@outlook.com

ARTICLE INFO


Keywords:

Chemical,
Residues,
Public Health,
Dysbiosis,
Phylogenics.

Vol.01, No.02, 2026

Copyright © 2026 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CCBY4.0).

<http://creativecommons.org/licenses/by/4.0/>



ABSTRACT

The escalating global demand for animal protein has driven the rapid intensification of livestock production, resulting in a heavy reliance on veterinary pharmaceuticals, agricultural pesticides, and intensive husbandry practices. While these interventions optimize growth and control disease, they introduce a critical food safety challenge: the persistence of chemical residues in edible animal tissues. This review synthesizes the primary classes of chemical contaminants including veterinary drugs, agrochemicals, heavy metals, and mycotoxins and traces their pathways into the human food chain. Chronic exposure to sub-lethal concentrations of these compounds poses significant consumer health risks, including the acceleration of global antimicrobial resistance (AMR), gut dysbiosis, carcinogenicity, and tissue-specific toxicity. Operational failures, such as non-adherence to mandatory drug withdrawal periods and weak regulatory oversight, remain the primary drivers of residue non-compliance. Ultimately, while post-harvest processing offers limited reduction of heat-labile compounds, securing food safety requires a fundamental shift toward farm-level mitigation, including strict analytical surveillance and the adoption of natural phytogetic and probiotic feed alternatives.

Introduction

The global trajectory of livestock production has shifted toward intensive, high-density confinement systems designed to maximize feed conversion ratios and satisfy the soaring worldwide demand for animal protein (Hernandez and Alagbe, 2025a). To sustain animal health and prevent catastrophic disease outbreaks within these crowded environments, modern animal husbandry relies heavily on a wide array of chemical interventions, ranging from prophylactic antimicrobials to synthetic growth promoters (Hernandez and Alagbe, 2025b). However, this systemic chemical dependency introduces a profound threat to food safety via the accumulation of chemical residues defined as the parent compounds, toxic metabolites, or associated impurities that persist within edible tissues like muscle, liver, kidney, and fat at the time of slaughter (Singh et al., 2022). This review provides a comprehensive synthesis of the primary chemical classes contaminating meat products, evaluates the complex entry pathways through which environmental and agricultural pollutants bioaccumulate in livestock, and establishes the specific physiological and toxicological mechanisms through which chronic, low-dose dietary exposure compromises human health (Adewale et al., 2021; Singh et al., 2021). By bridging the gap between field-level veterinary practice and clinical consumer pathology, this paper outlines the operational failures driving regulatory non-compliance and highlights sustainable, farm-level mitigation strategies necessary to secure a clean, chemical-safe global meat supply.

Entry Pathways of Chemical Residues

Veterinary Pharmaceuticals

Veterinary drugs are routinely administered to food-producing animals for therapeutic, metaphylactic, and prophylactic purposes, as well as for growth promotion (Madikizela et al., 2018). The most widely deployed pharmaceutical classes include broad-spectrum antimicrobials such as tetracyclines, sulfonamides, aminoglycosides (e.g., neomycin), and β -lactams (KumariAnjana, and

Jayachandran, 2017; Christou et al., 2017). When these compounds are misused, administered off-label, or given at inappropriate dosages, they fail to clear the animal's metabolic pathways prior to slaughter (Christou et al., 2017). Consequently, stable parent compounds and active metabolites remain bound to the skeletal muscle matrix or partition into highly vascularized internal organs, directly entering the human food supply (Adewole et al., 2016).

Agrochemical and Environmental Pollutants

Livestock frequently act as biological concentrators for persistent environmental pollutants present in contaminated pastures, forage, and drinking water (Bacanli, and Ba, 2019). Chief among these are organochlorine pesticides (OCPs) such as DDT, lindane, and aldrin, which enter the agricultural ecosystem through historical or current crop-protection practices (Donoghue, 2003). Because OCPs possess extreme chemical stability and high lipophilicity, they resist metabolic degradation within the animal. Instead, they readily cross biological membranes and progressively bioaccumulate within the adipose (fat) tissues of cattle, poultry, and rabbits over their production cycles (Donoghue, 2003).

Heavy Metals

The contamination of livestock tissue with toxic heavy metals such as lead (Pb), cadmium (Cd), and arsenic (As) is primarily driven by localized industrial emissions, mining runoff, and the application of contaminated wastewater or fertilizers to grazing lands (Manyi-Loh et al., 2018). Animals ingest these non-biodegradable elements directly from the soil while grazing or through contaminated commercial feed rations (Kimera et al., 2015). Once absorbed via the gastrointestinal tract, heavy metals mimic essential minerals and bind to systemic transport proteins, leading to a profound accumulation within chronic storage sites, most notably the renal cortex of the kidneys and the parenchyma of the liver ((Kimera et al., 2015).

Mycotoxins

Mycotoxins are toxic secondary metabolites produced by filamentous fungi, such as *Aspergillus*, *Penicillium*, and *Fusarium*, which routinely colonize cereal grains and oilseeds during pre-harvest growth or poor post-harvest storage. When livestock consume rations contaminated with Aflatoxin B1 (AFB1), the compound undergoes extensive cytochrome P450-mediated biotransformation in the liver. This metabolic pathway converts the toxin into its hydroxylated form, Aflatoxin M1 (AFM1), which retains high toxicity and carves out a persistent reservoir within the liver and kidney tissues of the animal, posing a direct threat to consumers who ingest offal (Nigeria et al., 2012).

Consumer Pathophysiology and Public Health Implications

Antimicrobial Resistance (AMR) & Gut Dysbiosis

The chronic ingestion of sub-therapeutic antimicrobial residues via meat consumption is a foundational driver of the global antimicrobial resistance crisis (Tavakoli et al., 2015). Continuous exposure to trace levels of antibiotics exerts selective evolutionary pressure within the consumer's gastrointestinal tract, killing sensitive commensal bacteria while allowing resistant strains to survive and proliferate (Er et al., 2013). This chronic exposure also disrupts the delicate taxonomic balance of the human intestinal microbiota, inducing a state of dysbiosis that compromises mucosal immunity, impairs nutrient absorption, and increases susceptibility to opportunistic enteric pathogens (Alagbe, 2026a,b).

Carcinogenicity, Genotoxicity, and Mutagenicity

A primary toxicological concern regarding long-term residue ingestion is the capacity of certain compounds to induce genetic mutations and malignant transformations. For instance, synthetic growth hormones like diethylstilbestrol (DES) act as potent endocrine disruptors that can induce reproductive organ cancers in humans (Salama et al., 2013). Similarly, organochlorine pesticides and feed-borne aflatoxin residues are recognized as highly genotoxic agents; they can bind directly to human DNA, forming covalent adducts that disrupt cellular replication and trigger hepatocarcinogenesis and tumor growth (Guetiya et al., 2016).

Target Organ Toxicity and Bioaccumulation Dynamics

Because different chemical contaminants exhibit distinct tissue affinities, their toxicological impacts are highly localized within specific human organ systems upon consumption (Alagbe et al., 2025). Heavy metals like cadmium and lead are cumulative toxins; chronic dietary intake causes them to pool in the human kidneys and bones, leading to irreversible nephropathy, renal tubular dysfunction, and bone demineralization (Alagbe, 2016). Meanwhile, residual aminoglycoside antibiotics concentrate

within the renal cortex and the endolymph of the inner ear, presenting distinct risks of sub-clinical nephrotoxicity and ototoxicity in vulnerable consumer populations (Alagbe, 2026).

Immunological Reactions

Beyond chronic toxicological pathways, chemical residues can trigger acute, immune-mediated hypersensitivity reactions in sensitized individuals. Trace residues of β -lactam antibiotics (such as penicillin) and sulfonamides present in meat products can act as haptens, binding to endogenous human proteins to elicit an allergic response (Muaz et al., 2018; Hammerum et al., 2007). These reactions can manifest clinically as mild dermatological eruptions, urticaria, or, in severe cases, life-threatening systemic anaphylactic shock (Van et al., 2015; Alberio et al., 2019).

Behavioral and Operational Drivers of Non-Compliance

The Economics of Withdrawal Periods

The leading operational cause of violative residue levels in meat is the failure of livestock producers to observe mandatory drug withdrawal periods the required timeframe between the final medication dose and the animal's slaughter (Yang et al., 2021). Under intense economic pressure, driven by high feed costs or fluctuating market prices, producers frequently rush livestock to market before the drug can be fully metabolized and excreted (Hiraku et al., 2004). This economic corner-cutting ensures that parent drug concentrations remain well above legal Maximum Residue Limits (MRLs) at the point of consumption.

Off-Label and Extra-Label Misuse

Residue compliance is heavily compromised by the widespread practice of extra-label drug use without adequate veterinary oversight (Huang et al., 2020). This includes modifying prescribed dosages, extending treatment durations, or administering pharmaceuticals to animal species for which they were never approved (Ahmed et al., 2020). Furthermore, the illicit use of completely banned growth-promoting agents, such as clenbuterol, creates highly volatile and dangerously elevated residue profiles in muscle tissues that completely bypass standard regulatory prediction models (Wu et al., 2024).

Feed Mill Cross-Contamination

Unintended chemical residues can also be traced back to technical errors within commercial feed manufacturing facilities (Mulchandani et al., 2021). When a single production line is sequentially utilized to blend medicated rations followed by standard finisher diets, mechanical carry-over and residual dust within mixing bins can easily cross-contaminate subsequent batches (Komarek et al., 2021; Abebe et al., 2021). This accidental introduction means that livestock can inadvertently ingest therapeutic agents right up until the day of slaughter, completely invalidating the farmer's intentional withdrawal schedules (Lees et al., 2021; Hosain et al., 2021).

Advanced Detection, Processing, and Farm-Level Mitigation Strategy

Analytical Instrumentation Frameworks

Robust regulatory surveillance and food safety enforcement depend entirely on structured, multi-tiered laboratory testing programs. Initial screening typically utilizes rapid, high-throughput diagnostic tools like Enzyme-Linked Immunosorbent Assay (ELISA) validation kits to identify suspect meat samples at the slaughterhouse. Positive matches are then subjected to definitive quantification using advanced analytical instrumentation, such as Atomic Absorption Spectroscopy (AAS) for heavy metal detection and Near-Infrared (NIR) feed analyzers to monitor incoming raw feed ingredients for contaminants before they enter the livestock metabolism.

Thermal and Processing Impacts

Relying on kitchen preparation to eliminate chemical risks is a dangerous strategy for consumers, as food processing methods yield highly variable results. While intensive thermal treatments such as prolonged boiling, deep-frying, or pressure cooking can thermally degrade or reduce the concentrations of certain heat-labile antibiotic molecules, they are completely ineffective against other threats (Patel et al., 2020). Toxic heavy metals (Pb, Cd) and highly stable environmental pollutants are completely heat-resistant; processing merely concentrates these non-volatile toxins within the cooked meat matrix as moisture is lost (Agren et al., 2025; Amenu et al., 2023).

Biosecurity and Green Alternatives

To achieve a sustainable reduction in chemical residues, livestock production must transition toward "green" farm-level alternatives that naturally reduce reliance on synthetic pharmaceuticals (Odey

et al., 2024). Integrating phytogenic feed additives, such as bioactive botanical extracts, provides natural antimicrobial and antioxidant support that improves animal performance without leaving harmful residues (Peng et al., 2024). Additionally, implementing strict on-farm biosecurity protocols, deploying homemade *Lactobacillus* probiotic serums, and utilizing solid-state fermented non-conventional feed resources can naturally boost livestock immunity and feed efficiency, minimizing the need for conventional veterinary drugs (Theodoridou et al., 2025; Kimera et al., 2020).

Conclusion

Chemical residue contamination in meat represents a multi-faceted threat to public health that requires a coordinated, "One Health" approach bridging veterinary medicine, environmental science, and human toxicology. As livestock systems continue to intensify, relying solely on end-product testing or post-harvest processing is insufficient to guarantee consumer safety. Future horizons must prioritize the digitization of livestock supply chains to enable real-time tracking of treatment histories, alongside the democratization of rapid, low-cost diagnostic screening tools at all processing points. Ultimately, safeguarding the consumer demands a structural shift toward sustainable animal nutrition and agro-ecological management, replacing routine chemical dependency with verified green alternatives to ensure long-term global food security.

References

- Hernandez, M and Alagbe, J.O. (2025). Influence of *Abrus procatorious* crude oil supplementation on the growth performance, Nutrient digestibility, Ruminal fermentation and Microbial population of Malabari Bucks. *International Journal of Global Sustainable Research*, 3(7): 527-538.
- Hernandez, M and Alagbe, J.O. (2025). Influence of *Odontonema strictum* oil on the growth performance and Ruminal fermentation of Barbari bucks. *Research in: Agriculture and Veterinary Sciences*, 9(2):1-10.
- Singh Sharma., Alagbe Olujimi John., Liu Xing., Sharma Ram and Kumar Amita (2022). Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectroemetry. *International Journal of Agriculture and Animal Production*, 2(6): 18-27.
- Adewale, A.O., Alagbe, J.O and Adeoye, K. (2021). Dietary Supplementation of *Rauvolfia vomitoria* Root extract as a Phytogenic Feed Additive in Growing Rabbit Diets: Haematology and Serum Biochemical Indices. *Concept in Dairy and Veterinary Sciences* 4(2): 2021.
- Singh, A.S., Alagbe, J.O., Sharma, S., Oluwafemi, R.A and Agubosi, O.C.P (2021). Effect of dietary supplementation of melon (*Citrallus linatus*) seed oil on the growth performance and oxidative status of growing rabbits. *Journal of Multidimensional Research and Reviews*, 2(1): 78-95.
- Madikizela, L.M.; Ncube, S.; Chimuka, L. Uptake of Pharmaceuticals by Plants Grown under Hydroponic Conditions and Natural Occurring Plant Species: A Review. *Sci. Total Environ.* 2018, 636, 477–486
- KumariAnjana, R.K.N.; Jayachandran, K.G.M.C. Persistence of Antibiotic Residue in Milk under Region of Bihar, India. *Int. J. Curr. Microbiol. Appl. Sci.* 2017, 6, 2296–2299.
- Christou, A.; Agüera, A.; Bayona, J.M.; Cytryn, E.; Fotopoulos, V.; Lambropoulou, D.; Manaia, C.M.; Michael, C.; Revitt, M.; Schröder, P.; et al. The Potential Implications of Reclaimed Wastewater Reuse for Irrigation on the Agricultural Environment: The Knowns and Unknowns of the Fate of Antibiotics and Antibiotic Resistant Bacteria and Resistance Genes—A Review. *Water Res.* 2017, 123, 448–467.
- Adewole, D.I.; Kim, I.H.; Nyachoti, C.M. Gut Health of Pigs: Challenge Models and Response Criteria with a Critical Analysis of the Effectiveness of Selected Feed Additives—A Review. *Asian-Australas. J. Anim. Sci.* 2016, 29, 909–924.
- Bacanli, M.; Ba,saran, N. Importance of Antibiotic Residues in Animal Food. *Food Chem. Toxicol.* 2019, 125, 462–466.
- Donoghue, D.J. Antibiotic Residues in Poultry Tissues and Eggs: Human Health Concerns? *Poult. Sci.* 2003, 82, 618–621.

Manyi-Loh, C.; Mamphweli, S.; Meyer, E.; Okoh, A. Antibiotic Use in Agriculture and Its Consequential Resistance in Environmental Sources: Potential Public Health Implications. *Molecules* 2018, 23, 795.

Kimera, Z.I.; Mdegela, R.H.; Mhaiki, C.J.N.; Karimuribo, E.D.; Mabiki, F.; Nonga, H.E.; Mwesongo, J. Determination of Oxytetracycline Residues in Cattle Meat Marketed in the Kilosa District, Tanzania. *Onderstepoort J. Vet. Res.* 2015, 82, 911.

Nigeria, I.A.; Olufemi, O.I.; Agboola, E.A. Oxytetracycline Residues in Edible Tissues of Cattle Slaughtered. *Niger. Vet. J.* 2011, 31, 2. 40. Bedada, A.H.; Zewde, B.M. Tetracycline Residue Levels in Slaughtered Beef Cattle from Three Slaughterhouses in Central Ethiopia. *Glob. Vet.* 2012, 8, 546–554.

Tavakoli, H.R.; Firouzabadi, M.S.S.; Afsharfarnia, S.; Jafari, N.J.; Sa'Adat, S. Detecting antibiotic residues by hplc method in chicken and calves meat in diet of a military center in tehran. *Acta Med. Mediterr.* 2015, 31, 1427–1433.

Er, B.; Onurdag, F.K.; Demirhan, B.; Ozgacar, S.Ö.; Oktem, A.B.; Abbasoglu, U. Screening of Quinolone Antibiotic Residues in Chicken Meat and Beef Sold in the Markets of Ankara, Turkey. *Poult. Sci.* 2013, 92, 2212–2215.

Salama, N.A.; Abou-Raya, S.H.; Shalaby, A.R.; Emam, W.H.; Mehaya, F.M. Incidence of Tetracycline Residues in Chicken Meat and Liver Retailed to Consumers. *Food Addit. Contam. Part B Surveill* 2011, 4, 88–93.

Guetiya Wadoun, R.E.; Zambou, N.F.; Anyangwe, F.F.; Njimou, J.R.; Coman, M.M.; Verdenelli, M.C.; Cecchini, C.; Silvi, S.; Orpianesi, C.; Cresci, A.; et al. Abusive Use of Antibiotics in Poultry Farming in Cameroon and the Public Health Implications. *Br. Poult. Sci.* 2016, 57, 483–493.

Muaz, K.; Riaz, M.; Akhtar, S.; Park, S.; Ismail, A. Antibiotic Residues in Chicken Meat: Global Prevalence, Threats, and Decontamination Strategies: A Review. *J. Food Prot.* 2018, 81, 619–627.

Hammerum, A.M.; Heuer, O.E.; Emborg, H.-D.; Bagger-Skjøt, L.; Jensen, V.F.; Rogues, A.-M.; Skov, R.L.; Agersø, Y.; Brandt, C.T.; Seyfarth, A.M.; et al. Danish Integrated Antimicrobial Resistance Monitoring and Research Program. *Emerg. Infect. Dis.* 2007, 13, 1633–1639.

Van Boeckel, T.P.; Brower, C.; Gilbert, M.; Grenfell, B.T.; Levin, S.A.; Robinson, T.P.; Teillant, A.; Laxminarayan, R. Global Trends in Antimicrobial Use in Food Animals. *Proc. Natl. Acad. Sci. USA* 2015, 112, 5649–5654.

Albero, B.; Tadeo, J.L.; Miguel, E.; Pérez, R.A. Rapid Determination of Antibiotic Residues in Cereals by Liquid Chromatography Triple Mass Spectrometry. *Anal. Bioanal. Chem.* 2019, 411, 6129–6139.

Hiraku, Y.; Sekine, A.; Nabeshi, H.; Midorikawa, K.; Murata, M.; Kumagai, Y.; Kawanishi, S. Mechanism of Carcinogenesis Induced by a Veterinary Antimicrobial Drug, Nitrofurazone, via Oxidative DNA Damage and Cell Proliferation. *Cancer Lett.* 2004, 215, 141–150.

Yang, X.; Chen, Z.; Zhao, W.; Liu, C.; Qian, X.; Zhang, M.; Wei, G.; Khan, E.; Hau Ng, Y.; Sik Ok, Y. Recent Advances in Photodegradation of Antibiotic Residues in Water. *Chem. Eng. J.* 2021, 405, 126806.

Huang, F.; An, Z.; Moran, M.J.; Liu, F. Recognition of Typical Antibiotic Residues in Environmental Media Related to Groundwater in China (2009–2019). *J. Hazard. Mater.* 2020, 399, 122813.

Ahmed, S.; Ning, J.; Peng, D.; Chen, T.; Ahmad, I.; Ali, A.; Lei, Z.; Abu bakr Shabbir, M.; Cheng, G.; Yuan, Z. Current Advances in Immunoassays for the Detection of Antibiotics Residues: A Review. *Food Agric. Immunol.* 2020, 31, 268–290.

Wu, D.; Dai, S.; Feng, H.; Karunaratne, S.; Yang, M.; Zhang, Y. Persistence and potential risks of tetracyclines and their transformation products in two typical different animal manure composting treatments. *Environ. Pollut.* 2024, 341, 122904.

Mulchandani, R.; Wang, Y.; Gilbert, M.; Van Boeckel, T.P. Global trends in antimicrobial use in food-producing animals: 2020 to 2030. *PLoS Glob. Public Health* 2023, 3, e0001305.

Komarek, A.M.; Dunston, S.; Enahoro, D.; Godfray, H.C.J.; Herrero, M.; Mason-D'Croz, D.; Rich, K.M.; Scarborough, P.; Springmann, M.; Sulser, T.B.; et al. Income, consumer preferences, and the future of livestock-derived food demand. *Glob. Environ. Change Hum. Policy Dimens.* 2021, 70, 102343.

- Abebe, E.; Gugsu, G.; Ahmed, M. Review on Major Food-Borne Zoonotic Bacterial Pathogens. *J. Trop. Med.* 2020, 2020, 4674235.
- Lees, P.; Pelligand, L.; Giraud, E.; Toutain, P.L. A history of antimicrobial drugs in animals: Evolution and revolution. *J. Vet. Pharmacol. Ther.* 2021, 44, 137–171.
- Hosain, M.Z.; Kabir, S.M.L.; Kamal, M.M. Antimicrobial uses for livestock production in developing countries. *Vet. World* 2021, 14, 210–221.
- Patel, S.J.; Wellington, M.; Shah, R.M.; Ferreira, M.J. Antibiotic Stewardship in Food-producing Animals: Challenges, Progress, and Opportunities. *Clin. Ther.* 2020, 42, 1649–1658.
- Agren, E.; Backhans, A.; Lindberg, M.; Sjolund, M.; Bengtsson, B.; Comin, A. Antimicrobial resistance in *Escherichia coli* from Swedish piglets with diarrhoea and associations with potential risk factors. *Acta Vet. Scand.* 2025, 67, 16.
- Amenu, K.; McIntyre, K.M.; Moje, N.; Knight-Jones, T.; Rushton, J.; Grace, D. Approaches for disease prioritization and decision-making in animal health, 2000-2021: A structured scoping review. *Front. Vet. Sci.* 2023, 10, 1231711.
- Odey, T.O.J.; Tanimowo, W.O.; Afolabi, K.O.; Jahid, I.K.; Reuben, R.C. Antimicrobial use and resistance in food animal production: Food safety and associated concerns in Sub-Saharan Africa. *Int. Microbiol. Off. J. Span. Soc. Microbiol.* 2024, 27, 1–23.
- Peng, C.; Ghanbari, M.; May, A.; Abeel, T. Effects of antibiotic growth promoter and its natural alternative on poultry cecum ecosystem: An integrated analysis of gut microbiota and host expression. *Front. Microbiol.* 2024, 15, 1492270.
- Theodoridou Oxinou, D.; Lamnisos, D.; Filippou, C.; Spervasilis, N.; Tsioutis, C. Antimicrobial Use and Antimicrobial Resistance in Food-Producing Animals: Cross-Sectional Study on Knowledge, Attitudes, and Practices Among Veterinarians and Operators of Establishments in the Republic of Cyprus. *Antibiotics* 2025, 14, 251.
- Kimera, Z.I.; Mshana, S.E.; Rweyemamu, M.M.; Mboera, L.E.G.; Matee, M.I.N. Antimicrobial use and resistance in food-producing animals and the environment: An African perspective. *Antimicrob. Resist. Infect. Control* 2020, 9, 37.
- Alagbe, J. O., Aliyu, K. I., Anaso, E. U., Anorue, D. N., & Ramalan, S. M. (2025). Impact of dietary exposure to heavy metal contaminated soil on the growth performance and hematological indices of growing rabbits. *Archives of Immunology Research and Therapy*, 5(1), 1–11.